



**You have downloaded a document from
RE-BUS
repository of the University of Silesia in Katowice**

Title: Application of multiple proxies in Mexican tropical coasts to prove evidence of tsunami deposits

Author: Krzysztof Gaidzik, María Teresa Ramírez-Herrera, Avto Goguitchaichvili, Francisco Bautista, Patricia Quintana, Ana-Carolina Ruiz-Fernández i in.

Citation style: Gaidzik Krzysztof, Ramírez-Herrera María Teresa, Goguitchaichvili Avto, Bautista Francisco, Quintana Patricia, Ruiz-Fernández Ana-Carolina i in. (2018). Application of multiple proxies in Mexican tropical coasts to prove evidence of tsunami deposits. "Geofísica internacional" Vol. 57, no 1 (2018), s. 9-10, doi 10.19155/geofint.2017.056.1.4



Uznanie autorstwa - Użycie niekomercyjne - Bez utworów zależnych Polska - Licencja ta zezwala na rozpowszechnianie, przedstawianie i wykonywanie utworu jedynie w celach niekomercyjnych oraz pod warunkiem zachowania go w oryginalnej postaci (nie tworzenia utworów zależnych).



UNIWERSYTET ŚLĄSKI
W KATOWICACH



Biblioteka
Uniwersytetu Śląskiego



Ministerstwo Nauki
i Szkolnictwa Wyższego

Application of multiple proxies in Mexican tropical coasts to prove evidence of tsunami deposits

Comment To: Ocampo-Rios, *et al.*, *Geofísica Internacional*, 56(1), 2017. DOI: 10.19155/geofint.2017.056.1.4

María Teresa Ramírez-Herrera*, Avto Goguitchaichvili, Francisco Bautista, Patricia Quintana, Ana-Carolina Ruiz-Fernández, Néstor Corona, Violeta Rangel, Marcelo Lagos, Vladimir Kostoglodov, María Luisa Machain, Daniel Aguilar Treviño, Rocío Castillo-Aja and Krzysztof Gaidzik

Received: August 17, 2017; accepted: October 06, 2017; published on line: January 01, 2018

The study of tsunami deposits has significantly advanced since the Chilean 2010 and Tohoku 2011 tsunamis providing opportunities to analyze tsunami deposits and their characteristics (Rubin *et al.*, 2017). In tropical environments, the combination of multiple proxies has demonstrated to be a necessity to prove evidence of ancient earthquakes and tsunamis (Ramírez-Herrera *et al.*, 2012, 2016; Williams *et al.*, 2011). Challenges faced in the study of tsunami deposits in tropical areas frequently affected by hurricanes, lead to problems of differentiation between tsunami and storm deposits, and misinterpretations of climate/seasonal changes.

Ocampo-Ríos *et al.* (2017) attempted to study the geologic evidence of the 1985 tsunami in Barra de Potosí, México. Their assertion that an "erosive base" is the only tool to prove the existence of tsunami deposits is not correct. (1) Our previous work in the Barra de Potosí area (field and survey-based interviews to witnesses of the 1985 tsunami) indicate that the area around the village has been intervened by human activity, thus surficial sediments are not reliable. (2) Beaches are very dynamic, and are located where normally tsunami erosion occur, thus, this type of environments are not suitable for tsunami deposits preservation. (3)

M.T. Ramírez-Herrera*
V. Rangel
Laboratorio Universitario de Geofísica Ambiental
Instituto de Geografía
Universidad Nacional Autónoma de México
*Corresponding author: tramirez@igg.unam.mx

A. Goguitchaichvili
Laboratorio Universitario de Geofísica Ambiental
Instituto de Geofísica
Universidad Nacional Autónoma de México
Unidad Michoacán, UNAM-Campus Morelia, México

F. Bautista
Laboratorio Universitario de Geofísica Ambiental
Centro de Investigaciones en Geografía Ambiental
Universidad Nacional Autónoma de México
Campus Morelia, Morelia, México

P. Quintana
D. Aguilar Treviño
Centro de Investigaciones y Estudios Avanzados,
Unidad Mérida, Yucatán, México

A.C. Ruiz-Fernández
Instituto de Ciencias del Mar y Limnología
Universidad Nacional Autónoma de México
Unidad Académica Mazatlán, Mazatlán, México

N. Corona
Centro de Estudios en Geografía Humana
El Colegio de Michoacán A.C., Michoacán, México

M. Lagos
Instituto de Geografía, Laboratorio de Tsunamis
Pontificia Universidad Católica de Chile
Santiago, Chile

V. Kostoglodov
Instituto de Geofísica
Universidad Nacional Autónoma de México
Ciudad de México, México

M.L. Machain
Instituto de Ciencias del Mar y Limnología
Universidad Nacional Autónoma de México
Ciudad Universitaria, Ciudad de México, México

R. Castillo-Aja
Posgrado en Geografía
Universidad Nacional Autónoma de México
Ciudad Universitaria, Ciudad de México, México

K. Gaidzik
Department of Fundamental Geology
University of Silesia, Poland

Ramírez-Herrera *et al.* (2012) research results come from sites in the Ixtapa estuary and not from Zihuatanejo Bay, i.e. a completely different geomorphic setting that consequently changes the tsunami impact and distribution of tsunami deposits. Thus, comparison by Ocampo-Ríos *et al.* (2017) with their sites is inadequate. (4) Ocampo-Ríos *et al.* (2017) hydraulic roughness calculation (0.02) to determine the inundation limit shows inaccuracies. Values from 0.011 to 0.02 apply to flood plains with very irregular shape which is not the case for Ixtapa estuary studied by Ramírez-Herrera *et al.* (2012) nor is the Zihuatanejo Bay. The calculation of Manning's values for the specific location (using the local topography, vegetation density, presence of barriers, etc.) is more appropriate than using standardized Manning's values. We reassessed here the tsunami flooding area interpreted by Ocampo-Ríos *et al.* (2017) using their data and demonstrate that their results are not correct, the inundation continues beyond 700 m in both Zihuatanejo and Barra de Potosí areas. (5) . Mineral content and assemblages are source-dependent and therefore, they are not a useful tool alone to identify tsunami deposits (Jagodziński *et al.*, 2012). (6) Except for the Br concentration values, Ocampo-Ríos *et al.* (2017) do not show significant differences in the elemental composition of the "pre-tsunami" and "tsunami deposits". The explanation provided on the low concentrations of Na, Cl and Br is not plausible. These elements have been widely used to identify marine provenance on sediment paleorecords along coastal areas. Br concentrations on soils can vary from 5 to 40 ppm, while on marine sediments they can reach up to 300 ppm (e.g. Ruiz-Fernández *et al.*, 2016). The oxides used to demonstrate tsunami origin of the Barra de Potosí sediments named "tsunami deposits", i.e. SiO₂ and TiO₂, if there was in fact any significant difference in values, would prove terrigenous characteristics and origin, and not marine. In summary, multiple proxies are required to prove evidence of tsunami deposits.

References

- Jagodziński R., Sternala B., Szczuciński W., Chagué-Goff C., Sugawara D., 2012, Heavy minerals in the 2011 Tohoku-oki tsunami deposits-insights into sediment sources and hydrodynamics. DOI:10.1016/j.sedgeo.2012.07.015
- Ocampo-Ríos B.G., D. Roy P., Macías M.C., Jonathan M.P., Lozano-Santacruz R., 2017, Tsunami deposits of September 21st 1985 in Barra de Potosí: comparison with other studies and evaluation of some geological proxies for southwestern Mexico. DOI:10.19155/geofint.2017.056.1.4
- Ramírez-Herrera M-T., Lagos M., Hutchinson I., Kostoglodov V., Machain M.L., Caballero M., Goguitchaichvili A., Aguilar B., Chagué-Goff C., Goff J., Ruiz-Fernández A.-C., Ortiz M., Nava H., Bautista F., López G.I., Quintana P., Extreme wave deposits on the Pacific coast of Mexico: Tsunamis or storms? — A multi-proxy approach. 2012. DOI:10.1016/j.geomorph.2011.11.002
- Ramírez-Herrera M.T., Bógalo M.-F., Černý J., Goguitchaichvili A., Corona N., Machain M.L., Carranza Edwards A., Sosa S., 2016, Historic and ancient tsunamis uncovered on the Jalisco-Colima Pacific coast, the Mexican subduction zone. *Geomorphology*, Elsevier, 2016, roč. 259, April. DOI:10.1016/j.geomorph.2016.02.011
- Ruiz-Fernández A.C., Sánchez-Cabeza J.-A., Serrato de la Peña J.L., Pérez-Bernal L.H., Cearreta A., Flores-Verdugo F., Machain-Castillo M.L., Chamizo E., García-Tenorio R., Queralt I., Dunbar R., Mucciarone D., Díaz-Asencio M., 2016, Accretion rates in coastal wetlands of the southeastern Gulf of California and their relationship with sea-level rise. DOI:10.1177/0959683616632882
- Rubín C.M., P. Horton B., Sieh K., Pilarczyk J.E., Daly P., Ismail N., Parnell A.C., 2017, Highly variable recurrence of tsunamis in the 7,400 years before the 2004 Indian Ocean tsunami. DOI:10.1038/ncomms16019.
- Williams S., Prasetya G., Chagué-Goff C., Goff J., Fai Cheung K., Davies T., Wilson T., 2011, Characterising diagnostic proxies for identifying palaeotsunamis in a tropical climatic regime, Samoan Islands. DOI:10.23919/OCEANS.2011.6107137